

Crystal structure of bis(μ_2 -thiocyanato)tetrakis(3,5-dimethyl-1-(thiocarbamoyl)pyrazole)dinickel(II) dichloride ethanol disolvate, $[\text{Ni}(\text{NCS})_2(\text{C}_6\text{H}_6\text{N}_3\text{S})_4][\text{Cl}]_2 \cdot 2\text{C}_2\text{H}_5\text{OH}$

Željko K. Jaćimović^{*I}, Ana Radović^{II}, Vukadin M. Leovac^{III}, Zoran D. Tomić^{IV} and Ivana Radosavljević Evans^V

^I University of Montenegro, Faculty of Metallurgy and Technology, 81000 Podgorica, Montenegro


^{II} Public Institution Railways of Montenegro, 81000 Podgorica, Montenegro

^{III} University of Novi Sad, Faculty of Sciences, Institute of Chemistry, Trg Dositeja Obradovića 3, 21000 Novi Sad, Serbia

^{IV} Vinča Institute of Nuclear Sciences, Laboratory of Theoretical Physics and Condensed Matter Physics, 11001 Belgrade, Serbia

^V University of Durham, Department of Chemistry, South Road, Durham DH1 3LE, England

Received August 8, 2007; accepted and available on-line January 7, 2008; CCDC no. 1267/2162



rameters of the two chelate rings differ significantly from those observed in square-planar Ni(L₂)₂ complex [10] and indicate weaker bonding. The bridging thiocyanate group is nearly linear and shows bent coordination modes with the metal atoms ($\angle \text{S3}–\text{C22}–\text{N11} = 178.58^\circ$, $\angle \text{Ni1}–\text{S3}–\text{C22} = 102.40^\circ$, $\angle \text{Ni1}–\text{N11}–\text{C22} = 164.47^\circ$). Distance between the two nickel atoms is 5.578 Å. The packing of complex molecules is determined by the presence of free chloride anion, which ethanol molecule. Repulsion of the chloride anion lead to the arrangement of hydrogen bonded atoms in a tetrahedral fashion. The other impor-

tant factor is the position of hydrogen bond donors in the complex molecule. Both hydrogen atoms at nitrogen N14 are involved in the hydrogen bonds with Cl, and formation of molecular chain, while the connection between the neighboring chains is achieved by the hydrogen bond involving atom N13 and chlorine. The angle between the two neighboring molecular chains is 69° (measured by the torsion angle formed by the lines connecting the centroids of the metallocycle rings). The crystal structure is additionally stabilized by the hydrogen bond involving ethanol molecule and nitrogen N13.

Table 1. Data collection and handling.

Crystal:	green prism, size 0.04 × 0.16 × 0.20 mm
Wavelength:	Mo K α radiation (0.71073 Å)
μ :	12.82 cm ⁻¹
Diffractometer, scan mode:	Bruker AXS SMART CCD, ω/ϕ
2 θ_{max} :	59.98°
$N(hkl)_{\text{measured}}$, $N(hkl)_{\text{unique}}$:	27632, 6381
Criterion for I_{obs} , $N(hkl)_{\text{gt}}$:	$I_{\text{obs}} > 2 \sigma(I_{\text{obs}})$, 4062
$N(\text{param})_{\text{refined}}$:	349
Programs:	SHELXS-97 [12], SHELXL-97 [13], ORTEP-3 [14], WinGX [15], ORTEP-II [16]

Table 2. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso}
H(5)	4e	0.661(2)	0.486(1)	−0.014(1)	0.040(2)
H(6)	4e	0.630(2)	0.463(1)	0.075(1)	0.041(2)
H(7)	4e	0.562(2)	0.062(1)	0.234(1)	0.033(2)
H(8)	4e	0.318(2)	0.128(3)	−0.001(1)	0.051(2)

Table 2. Continued.

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso}
H(9)	4e	0.452(2)	0.1934(9)	−0.118(1)	0.037(2)
H(10)	4e	0.751(2)	0.170(2)	−0.155(2)	0.056(2)
H(11)	4e	0.379(2)	0.062(2)	0.075(2)	0.051(2)
H(12)	4e	0.769(2)	0.598(2)	−0.099(2)	0.068(2)
H(14)	4e	0.487(1)	0.136(1)	0.180(1)	0.034(2)
H(15)	4e	0.362(2)	0.190(2)	0.081(2)	0.051(2)
H(16)	4e	0.986(1)	0.447(2)	−0.1622(9)	0.051(2)
H(17)	4e	0.895(2)	0.630(2)	−0.092(2)	0.068(2)
H(18)	4e	0.638(2)	0.185(2)	−0.207(1)	0.057(2)
H(22)	4e	1.060(3)	0.166(2)	−0.084(1)	0.084(2)
H(23)	4e	0.834(3)	0.610(2)	−0.010(1)	0.069(2)
H(25)	4e	0.976(2)	0.158(2)	−0.163(2)	0.084(2)
H(30)	4e	1.076(2)	0.239(2)	−0.164(2)	0.084(2)
H(34)	4e	0.695(2)	0.288(2)	−0.161(2)	0.057(2)
H(1)	4e	0.319(3)	0.384(2)	−0.163(2)	0.099(2)
H(2)	4e	0.196(2)	0.397(2)	−0.200(2)	0.098(2)
H(3)	4e	0.232(2)	0.261(2)	−0.303(1)	0.052(2)
H(4)	4e	0.401(3)	0.187(2)	−0.295(2)	0.051(2)
H(13)	4e	0.257(2)	0.210(2)	−0.211(1)	0.052(2)
H(241)	4e	0.296(3)	0.436(2)	−0.253(2)	0.099(2)

Table 3. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> ₁₁	<i>U</i> ₂₂	<i>U</i> ₃₃	<i>U</i> ₁₂	<i>U</i> ₁₃	<i>U</i> ₂₃
Ni(1)	4e	0.83558(3)	0.15012(3)	0.04438(2)	0.0158(1)	0.0200(2)	0.0178(1)	0.0011(1)	0.0042(1)	0.0010(1)
S(2)	4e	0.74876(5)	0.03482(6)	0.14810(4)	0.0182(3)	0.0267(3)	0.0200(3)	0.0018(2)	0.0019(2)	0.0055(2)
S(3)	4e	1.01656(5)	0.13779(6)	0.12758(4)	0.0179(3)	0.0264(3)	0.0260(3)	0.0009(2)	0.0015(2)	−0.0067(3)
S(5)	4e	0.79333(6)	0.31496(6)	0.13741(4)	0.0297(3)	0.0252(3)	0.0211(3)	0.0041(3)	0.0054(2)	0.0002(2)
N(8)	4e	0.5896(2)	0.1382(2)	0.0490(1)	0.0161(9)	0.023(1)	0.0151(9)	0.0011(8)	0.0037(7)	−0.0001(8)
N(9)	4e	0.8347(2)	0.3830(2)	−0.0187(1)	0.026(1)	0.021(1)	0.024(1)	0.0006(9)	0.0052(9)	0.0034(8)
N(10)	4e	0.6724(2)	0.1582(2)	−0.0058(1)	0.0180(9)	0.025(1)	0.0144(9)	0.0021(8)	0.0052(7)	0.0017(8)
N(11)	4e	0.8717(2)	0.0255(2)	−0.0362(1)	0.0164(9)	0.028(1)	0.023(1)	0.0004(9)	0.0021(8)	−0.0012(9)
N(12)	4e	0.8873(2)	0.2819(2)	−0.0309(1)	0.021(1)	0.024(1)	0.026(1)	0.0020(8)	0.0075(8)	0.0022(9)
N(13)	4e	0.5472(2)	0.0962(2)	0.1867(1)	0.021(1)	0.031(1)	0.019(1)	0.0017(9)	0.0053(8)	0.0035(9)
N(14)	4e	0.6762(2)	0.4548(2)	0.0350(2)	0.030(1)	0.031(1)	0.026(1)	0.008(1)	0.0052(9)	0.001(1)
C(1)	4e	1.0223(3)	0.2089(3)	−0.1286(2)	0.044(2)	0.045(2)	0.053(2)	0.007(2)	0.036(2)	0.013(2)
C(7)	4e	0.8717(2)	0.4652(2)	−0.0717(2)	0.025(1)	0.027(1)	0.035(2)	−0.005(1)	0.002(1)	0.011(1)
C(15)	4e	0.9471(2)	0.4136(3)	−0.1196(2)	0.027(1)	0.040(2)	0.040(2)	−0.002(1)	0.010(1)	0.018(1)
C(16)	4e	0.6210(2)	0.0923(2)	0.1295(2)	0.021(1)	0.019(1)	0.016(1)	−0.0029(9)	0.0018(8)	−0.0003(9)
C(17)	4e	0.5041(2)	0.1791(2)	−0.0748(2)	0.025(1)	0.035(2)	0.018(1)	0.005(1)	−0.0001(9)	0.000(1)
C(18)	4e	0.7635(2)	0.3903(2)	0.0489(2)	0.026(1)	0.020(1)	0.024(1)	−0.000(1)	0.003(1)	−0.004(1)
C(19)	4e	0.9535(2)	0.3001(3)	−0.0939(2)	0.025(1)	0.038(2)	0.030(1)	0.001(1)	0.012(1)	0.009(1)
C(20)	4e	0.4853(2)	0.1500(2)	0.0065(2)	0.018(1)	0.030(1)	0.019(1)	0.003(1)	−0.0014(8)	−0.004(1)
C(21)	4e	0.3763(2)	0.1303(3)	0.0432(2)	0.016(1)	0.046(2)	0.024(1)	0.000(1)	0.0005(9)	−0.005(1)
C(22)	4e	1.0834(2)	0.0425(2)	0.0744(2)	0.015(1)	0.026(1)	0.020(1)	−0.0031(9)	−0.0001(8)	0.003(1)
C(23)	4e	0.6205(2)	0.1832(2)	−0.0802(2)	0.024(1)	0.028(1)	0.016(1)	0.005(1)	0.0014(9)	0.0009(9)
C(25)	4e	0.6816(3)	0.2091(3)	−0.1578(2)	0.032(1)	0.044(2)	0.019(1)	0.004(1)	0.005(1)	0.006(1)
C(27)	4e	0.8388(3)	0.5859(3)	−0.0673(2)	0.027(1)	0.030(2)	0.059(2)	−0.006(1)	−0.002(1)	0.017(2)
O(6)	4e	0.3921(2)	0.2465(2)	−0.2682(1)	0.027(1)	0.029(1)	0.030(1)	0.0005(8)	0.0032(8)	0.0013(8)
C(24)	4e	0.2717(3)	0.3805(3)	−0.2140(3)	0.036(2)	0.052(2)	0.079(3)	−0.005(2)	0.013(2)	−0.035(2)
C(26)	4e	0.2792(2)	0.2670(3)	−0.2505(2)	0.023(1)	0.035(2)	0.050(2)	−0.003(1)	0.003(1)	−0.005(1)
Cl(4)	4e	0.54856(6)	−0.03139(6)	0.36392(4)	0.0274(3)	0.0378(4)	0.0213(3)	0.0030(3)	0.0042(2)	0.0022(3)

References

1. Radosavljević Evans, I.; Howard, J. A. K.; Howard, L. E. M.; Evans, J. S. O.; Jaćimović, Ž. K.; Jevtović, V. S.; Leovac, V. M.: Transition metal complexes with pyrazole based ligands, part 18: new binuclear Cu(I), Cu(II) and Co(II) complexes with 3,5-dimethyl-1-thiocarboxamide pyrazole: synthesis, structural and magnetic studies. *Inorg. Chim. Acta* **357** (2004) 4528-4536.
2. Radosavljević Evans, I.; Mészáros Szécsényi, K.; Leovac, V. M.: Di- μ -chloro-bis{chloro[3,5-dimethyl-1-(thiocarbamoyl)pyrazole- $\kappa^2 N^2, S$]cadmium(II)}. *Acta Crystallogr. E* **61** (2005) m641-m643.
3. Tomić, Z. D.; Jaćimović, Ž. K.; Leovac, V. M.; Cesljević, V. I.: Transition metal complexes with pyrazole-derived ligands. XI. [Zn(μ -L)(HL)(OAc)]₂ (HL is 4-acetyl-3-amino-5-methylpyrazole). *Acta Crystallogr. C* **56** (2000) 777-779.
4. Kovács, A.; Nemcsok, D.; Pokol, G.; Mészáros Szécsényi, K.; Leovac, V. M.; Jaćimović, Ž. K.; Radosavljević Evans, I.; Howard, J. A. K.; Tomić, Z. D.; Giester, G.: Structural, spectroscopic and computational studies of the HgL₂Cl₂ complex (L = 3,5-dimethyl-1-thiocarboxamide pyrazole) and the crystal structure of L. *New J. Chem.* **29** (2005) 833-840.
5. Nemcsok, D.; Kovács, A.; Mészáros Szécsényi, K.; Leovac, V. M.: Vibrational spectroscopic and theoretical study of 3,5-dimethyl-1-thiocarboxamide pyrazole (L) and the complexes Co₂L₂Cl₄, Cu₂L₂Cl₄ and Cu₂L₂Br₂. *Chem. Phys.* **328** (2006) 85-92.
6. Barik, A. K.; Paul, S.; Butcher, R. J.; Kar, S. K.: Synthesis and characterization of two tris-chelate complexes of cobalt(III) with 3,5-dimethyl-1-(N-methyl/ethyl)thiocarbamylpyrazole (HL1, HL2) – biologically important bidentate ligands with one ambidentate donor site. *Polyhedron* **19** (2000) 2651-2655.
7. Jaćimović, Ž. K.; Tomić, Z. D.; Giester, G.; Leovac, V. M.: Tris(3,5-dimethyl-1H-pyrazole-1-thiocarboxamidato- $\kappa^2 N^2, N$)cobalt(III). *Acta Crystallogr. C* **59** (2003) m381-m383.
8. Pearson, R. G.: Hard and Soft Acids and Bases. *J. Am. Chem. Soc.* **85** (1963) 3533-3539.
9. Barik, A. K.; Bandyopadhyay, P.; Kar, S. K.: Copper(II) and nickel(II) complexes of pyrazole derived ligands: Synthesis, characterization and coordinating properties of two substituted thiocarbamyl pyrazoles, 3,5-dimethyl-1-N-methyl/ethyl thiocarbamyl pyrazole (HL1, HL2), potential ligands for biological interest. X-ray crystallographic studies of Ni(L₂)₂. *Polyhedron* **18** (1999) 1-6.
10. Radosavljević Evans, I.; Howard, J. A. K.; Mészáros Szécsényi, K.; Leovac, V. M.; Jaćimović, Ž. K.: Synthesis, Characterization and Crystal Structure of a Novel Ni(II) Complex, Ni(L-H)₂ (L = 3,5-dimethyl-1-thiocarboxamide pyrazole). *J. Coord. Chem.* **57** (2004) 469-476.
11. Sheldrick, G. M.: SHELXS-97. Program for the Solution of Crystal Structures. University of Göttingen, Germany 1997.
12. Sheldrick, G. M.: SHELXL-97. Program for the Refinement of Crystal Structures. University of Göttingen, Germany 1997.
13. Farrugia, L. J.: ORTEP-3 for Windows – a version of ORTEP-III with a Graphical User Interface. *J. Appl. Crystallogr.* **30** (1997) 565.
14. Farrugia, L. J.: WinGX suite for small-molecule single-crystal crystallography. *J. Appl. Crystallogr.* **32** (1999) 837-838.
15. Johnson, C. K.: ORTEP-II. A Fortran Thermal-Ellipsoid Plot Program. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA 1976.